

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 38/16		A1	(11) International Publication Number: WO 00/12116
			(43) International Publication Date: 9 March 2000 (09.03.00)
(21) International Application Number: PCT/US99/19348			(81) Designated States: AE, AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CR, CU, CZ, DM, EE, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, RO, RU, SD, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
(22) International Filing Date: 24 August 1999 (24.08.99)			
(30) Priority Data: 60/098,273 28 August 1998 (28.08.98) US 60/100,012 11 September 1998 (11.09.98) US			
(71) Applicant (for all designated States except US): ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): HUGHES, Benjamin, Lee [US/US]; 402 Blue Ridge Road, Indianapolis, IN 46208 (US). WOLFF, Ronald, Keith [US/US]; 12329 Brookshire Parkway, Carmel, IN 46033 (US).			
(74) Agents: MACIAK, Ronald, S. et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).			Published With international search report.
(54) Title: METHOD FOR ADMINISTERING INSULINOTROPIC PEPTIDES			
(57) Abstract <p>The claimed invention relates to a method of administering glucagon-like peptide-1 molecules by inhalation, a method for treating diabetes by administering glucagon-like peptide-1 molecules by inhalation, and a method for treating hyperglycemia by administering glucagon-like peptide-1 molecules by inhalation.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-1-

METHOD FOR ADMINISTERING INSULINOTROPIC PEPTIDES

Field of the Invention

This invention relates to methods of treating
5 humans suffering from diabetes and insulin resistance. In particular, the invention relates to the pulmonary delivery of glucagon-like peptide-1 (GLP-1) and analogs thereof for systemic absorption through the lungs to eliminate the need for administering anti-diabetic compounds by injection.

10 Background of the Invention

Glucagon-like peptide-1 was first identified in 1987 as a incretin hormone, a peptide secreted by the gut upon ingestion of food. Glucagon-like peptide-1 is secreted by the L-cells of the intestine after being proteolytically
15 processed from the 160 amino acid precursor protein, preproglucagon. Cleavage of preproglucagon first yields glucagon-like peptide-1, a 37 amino acid peptide that is poorly active. A subsequent cleavage of the peptide bond between residues 6 and 7 yields biologically active
20 glucagon-like peptide-1 referred to as GLP-1(7-37). It should be noted that this specification uses the nomenclature scheme that has developed around this hormone. By convention in the art, the amino terminus of GLP-1(7-37) has been assigned number 7 and the carboxy terminus number
25 37. Approximately 80% of the GLP-1(7-37) that is synthesized is amidated at the C-terminal after removal of the terminal glycine residue in the L-cells. The biological effects and metabolic turnover of the free acid GLP-1(7-37), and the amide, GLP-1(7-36)NH₂, are indistinguishable. As

-2-

used herein, these two naturally-occurring forms will be referred to collectively as GLP-1.

GLP-1 is known to stimulate insulin secretion (insulinotropic action) causing glucose uptake by cells which decreases serum glucose levels (see, e.g., Mojsov, S., *Int. J. Peptide Protein Research*, 40:333-343 (1992)). Numerous GLP-1 analogs and derivatives demonstrating insulinotropic action are known in the art. Also it has been demonstrated that the N-terminal histidine residue (His 7) is very important to insulinotropic activity of GLP-1 (Suzuki, S., et al. *Diabetes Res.; Clinical Practice* 5 (Supp. 1):S30 (1988)).

Multiple authors have demonstrated the nexus between laboratory experimentation and mammalian, particularly human, insulinotropic responses to exogenous administration of GLP-1. See, e.g., Nauck, M.A., et al., *Diabetologia*, 36:741-744 (1993); Gutniak, M., et al., *New England J. of Medicine*, 326(20):1316-1322 (1992); Nauck, M.A., et al., *J. Clin. Invest.*, 91:301-307 (1993); and Thorens, B., et al., *Diabetes*, 42:1219-1225 (1993)].

GLP-1 based peptides hold great promise as alternatives to insulin therapy for patients with diabetes who have failed on sulfonylureas. GLP-1 has been studied intensively by academic investigators, and this research has established the following for patients with type II diabetes who have failed on sulfonylureas:

1) GLP-1 stimulates insulin secretion, but only during periods of hyperglycemia. The safety of GLP-1 compared to insulin is enhanced by this property of GLP-1 and by the observation that the amount of insulin secreted is

-3-

proportional to the magnitude of the hyperglycemia. In addition, GLP-1 therapy will result in pancreatic release of insulin and first-pass insulin action at the liver. This results in lower circulating levels of insulin in the periphery compared to subcutaneous insulin injections.

2) GLP-1 suppresses glucagon secretion, and this, in addition to the delivery of insulin via the portal vein helps suppress the excessive hepatic glucose output in diabetic patients.

3) GLP-1 slows gastric emptying which is desirable in that it spreads nutrient absorption over a longer time period, decreasing the postprandial glucose peak.

4) Several reports have suggested that GLP-1 may enhance insulin sensitivity in peripheral tissues such as muscle and fat.

5) Finally, GLP-1 has been shown to be a potential regulator of appetite.

Meal-time use of GLP-1 based peptides offers several advantages over insulin therapy. Insulin therapy requires blood glucose monitoring, which is both expensive and painful. The glucose-dependency of GLP-1 provides an enhanced therapeutic window in comparison to insulin, and should minimize the need to monitor blood glucose. Weight gain also can be a problem with intensive insulin therapy, particularly in the obese type II diabetic patients.

The therapeutic potential for native GLP-1 is further increased if one considers its use in patients with type I diabetes. A number of studies have demonstrated the effectiveness of native GLP-1 in the treatment of insulin dependent diabetes mellitus. Similar to patients with type

-4-

II diabetes, GLP-1 is effective in reducing fasting hyperglycemia through its glucagonostatic properties. Additional studies have indicated that GLP-1 also reduces postprandial glycemic excursions in type I patients, most likely through a delay in gastric emptying. These observations indicate that GLP-1 may be useful as a treatment in type I and type II patients.

To date administration of clinically proven peptide hormones and as well as GLP-1 has generally been accomplished by subcutaneous injection which is both inconvenient and unattractive. Therefore, many investigators have studied alternate routes for administering peptide hormones such as oral, rectal, transdermal, and nasal routes. Thus far, however, these routes of administration have not resulted in clinically proven peptide hormone therapy.

It has been known for a number of years that some proteins can be absorbed from the lung. For example, insulin administered by inhalation aerosol to the lung was first reported by Gaensslen in 1925. Despite the fact that a number of human and animal studies have shown that some insulin formulations can be absorbed through the lungs, pulmonary delivery of peptide hormones has not been vigorously pursued because of very low bioavailability. Larger proteins, such as cytokines and growth factors which are generally larger than 150 amino acid residues, are often readily absorbed by the cells lining the alveolar regions of the lung. Pulmonary absorption of smaller proteins is however much less predictable; though insulin (51 residues), calcitonin (32 residues) and parathyroid hormone

-5-

(34 residues) have been reported to be systemically absorbed through the pulmonary route. See US Patent No: 5,607,915, herein incorporated by reference. Despite systemic absorption by the lung of some small protein hormones, the pharmacodynamics associated with pulmonary delivery of peptides is unpredictable.

Thus, there is a need to provide a reliable pulmonary method of delivering GLP-1 and related analogs because it would offer patients an attractive, non-invasive alternative to insulin. This need is particularly true since insulin has a very narrow therapeutic index while GLP-1 treatment offers a way to normalize blood glucose only in response to hyperglycemic conditions without the threat of hypoglycemia.

Not all protein hormones can be efficiently absorbed through the lungs, and there are many factors that affect it. Absorption of proteins in the lung is largely dependent on the physical characteristics of the protein. Thus, even though pulmonary delivery of some protein hormones has been observed, the physical properties and short length of GLP-1 and some related peptides made it unclear whether such peptides could be effectively delivered through the pulmonary route.

Efficient pulmonary delivery is dependent on the ability to deliver the protein to the deep lung alveolar epithelium. Protein particles that lodge in the upper airway epithelium are not absorbed to a significant extent because the overlying mucus functions to trap, and then clear debris by mucociliary transport up the airway. This mechanism is also a major contributor to low

-6-

bioavailability. The extent to which proteins are not absorbed and instead eliminated by these routes depends on their solubility, their size, and other largely uncharacterized mechanisms.

5 Even when a peptide hormone can be reproducibly delivered to the deep lung alveolar epithelium, it is difficult to predict whether it will be rapidly absorbed and transported to the blood. Absorption values for some proteins delivered through the lungs have been calculated
10 and range from fifteen minutes for parathyroid hormone (1-34) to 48 hours for glycosylated α 1-antitrypsin. Moreover a variety of endogenous peptidases exist in the lung which can degrade peptides prior to absorption. Thus, the longer it takes for a peptide particle to dissolve and be absorbed,
15 the greater the chance for enzymatic inactivation. Thus, because of the small size of GLP-1 and its inherent susceptibility to certain enzymes, it was most surprising to find that an aerosolized GLP-1 analog could be reproducibly and effectively delivered through the lungs.

20 Summary of the Invention

 The present invention relates to a method for administering a glucagon-like peptide-1 molecule comprising, administering an effective amount of the peptide to a patient in need thereof by pulmonary delivery. The present
25 invention also relates to a method for treating diabetes comprising, administering an effective dose of a glucagon-like peptide-1 to a patient in need thereof by pulmonary delivery. Another aspect of the invention relates to a method for treating hyperglycemia comprising, administering
30 an effective dose of a glucagon-like peptide-1 to a patient

-7-

in need thereof by pulmonary delivery. Preferably, the glucagon-like peptide-1 molecule is delivered by inhalation and to the lower airway of the patient.

The glucagon-like peptide-1 can be delivered in a carrier, as a solution or suspension, or as a dry powder, using any of a variety of devices suitable for administration by inhalation. Preferably, the glucagon-like peptide-1 is delivered in a particle size effective for reaching the lower airways of the lung.

10 Detailed Description of the Invention

The term "GLP-1" refers to human glucagon-like peptide-1 whose sequences and structures are known in the art. See US Patent No. 5,120,712, herein incorporated by reference. As previously discussed, there are two native forms of human GLP-1, GLP-1(7-37) and GLP-1(7-36)NH₂, which will be distinguished only when necessary.

The term "GLP-1 analog" is defined as a molecule having one or more amino acid substitutions, deletions, inversions, or additions compared with GLP-1. Many GLP-1 analogs are known in the art and include, for example, GLP-1(7-34) and GLP-1(7-35), GLP-1(7-36), Val⁸-GLP-1(7-37), Gln⁹-GLP-1(7-37), D-Gln⁹-GLP-1(7-37), Thr¹⁶-Lys¹⁸-GLP-1(7-37), and Lys¹⁸-GLP-1(7-37). Preferred GLP-1 analogs are GLP-1(7-34) and GLP-1(7-35), which are disclosed in U.S. Patent No: 5,118,666, herein incorporated by reference.


The term "GLP-1 derivative" is defined as a molecule having the amino acid sequence of GLP-1 or a GLP-1 analog, but additionally having chemical modification of one or more of its amino acid side groups, a-carbon atoms, terminal amino group, or terminal carboxylic acid group. A

-8-

chemical modification includes, but is not limited to, adding chemical moieties, creating new bonds, and removing chemical moieties. Modifications at amino acid side groups include, without limitation, acylation of lysine e-amino groups, N-alkylation of arginine, histidine, or lysine, alkylation of glutamic or aspartic carboxylic acid groups, and deamidation of glutamine or asparagine. Modifications of the terminal amino include, without limitation, the des-amino, N-lower alkyl, N-di-lower alkyl, and N-acyl modifications. Modifications of the terminal carboxy group include, without limitation, the amide, lower alkyl amide, dialkyl amide, and lower alkyl ester modifications. Lower alkyl is C₁-C₄ alkyl. Furthermore, one or more side groups, or terminal groups, may be protected by protective groups known to the ordinarily-skilled protein chemist. The α -carbon of an amino acid may be mono- or di-methylated.

The term "GLP-1 molecule" means GLP-1, GLP-1 analog, or GLP-1 derivative.

Another preferred group of GLP-1 analogs is defined by the formula:



and the pharmaceutically-acceptable salts thereof, wherein: R₁ is selected from the group consisting of L-histidine, D-histidine, desamino-histidine, 2-amino-histidine, β -hydroxy-histidine, homohistidine, α -fluoromethyl-histidine, and α -methyl-histidine; X is selected from the group consisting of Ala, Gly, Val, Thr, Ile, and α -methyl-Ala;

-9-

Y is selected from the group consisting of Glu, Gln, Ala, Thr, Ser, and Gly; Z is selected from the group consisting of Glu, Gln, Ala, Thr, Ser, and Gly; and R₂ is selected from the group consisting of NH₂, and Gly-OH; providing that when
5 R₁ is His, X is Ala, Y is Glu, and Z is Glu, R₂ must be NH₂.

Yet another preferred group of compounds consistent with the present invention is disclosed in WO 91/11457 (U.S. Patent No. 5,545,618, herein incorporated by reference) and consists essentially of GLP-1(7-34), GLP-1(7-
10 35), GLP-1(7-36), or GLP-1(7-37), or the amide forms thereof, and pharmaceutically-acceptable salts thereof, having at least one modification selected from the group consisting of:

(a) substitution of glycine, serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine, isoleucine, leucine, methionine, phenylalanine, arginine, or D-lysine for lysine at position 26 and/or position 34; or substitution of glycine, serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine, isoleucine, leucine, methionine, phenylalanine, lysine, or a
20 D-arginine for arginine at position 36;

(b) substitution of an oxidation-resistant amino acid for tryptophan at position 31;

(c) substitution of at least one of: tyrosine for valine at position 16; lysine for serine at position 18; aspartic acid for glutamic acid at position 21; serine for glycine at position 22; arginine for glutamine at position 23; arginine for alanine at position 24; and glutamine for lysine at position 26; and
25

-10-

(d) substitution of at least one of: glycine, serine, or cysteine for alanine at position 8; aspartic acid, glycine, serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine, isoleucine, leucine, methionine, or phenylalanine for glutamic acid at position 9; serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine, isoleucine, leucine, methionine, or phenylalanine for glycine at position 10; and glutamic acid for aspartic acid at position 15; and

(e) substitution of glycine, serine, cysteine, threonine, asparagine, glutamine, tyrosine, alanine, valine, isoleucine, leucine, methionine, or phenylalanine, or the D- or N-acylated or alkylated form of histidine for histidine at position 7; wherein, in the substitutions is (a), (b), (d), and (e), the substituted amino acids can optionally be in the D-form and the amino acids substituted at position 7 can optionally be in the N-acylated or N-alkylated form.

Because the enzyme, dipeptidyl-peptidase IV (DPP IV), may be responsible for the observed rapid *in vivo* inactivation of administered GLP-1, [see, e.g., Mentlein, R., et al., *Eur. J. Biochem.*, 214:829-835 (1993)], administration of GLP-1 analogs and derivatives that are protected from the activity of DPP IV is preferred, and the administration of Gly⁸-GLP-1(7-36)NH₂, Val⁸-GLP-1(7-37)OH, a-methyl-Ala⁸-GLP-1(7-36)NH₂, and Gly⁸-Gln²¹-GLP-1(7-37)OH, or pharmaceutically-acceptable salts thereof, is more preferred.

The use in the present invention of a molecule claimed in U.S. Patent No. 5,188,666, herein incorporated by reference, is preferred. Such molecule is selected from the

-11-

group consisting of a peptide having the amino acid sequence:

His-Ala-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-Ser-Tyr-
Leu-Glu-Gly-Gln-Ala-Ala-Lys-Glu-Phe-Ile-Ala-Trp-Leu-
5 Val-X (SEQ ID NO:2)

wherein X is selected from the group consisting of Lys and Lys-Gly; and a derivative of said peptide, wherein said peptide is selected from the group consisting of: a pharmaceutically-acceptable acid addition salt of said
10 peptide; a pharmaceutically-acceptable carboxylate salt of said peptide; a pharmaceutically-acceptable lower alkylester of said peptide; and a pharmaceutically-acceptable amide of said peptide selected from the group consisting of amide, lower alkyl amide, and lower dialkyl amide.

15 Another preferred group of molecules for use in the present invention consists of compounds disclosed in U.S. Patent No. 5,512,549, herein incorporated by reference, having the general formula:

R¹-Ala-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-Ser-Tyr-
20 Leu-Glu-Gly-Gln-Ala-Ala-Xaa-Glu-Phe-Ile-Ala-Trp-
Leu-Val-Lys-Gly-Arg-R³ R²

(SEQ ID NO:3)

and pharmaceutically-acceptable salts thereof, wherein R¹ is
25 selected from the group consisting of 4-imidazopropionyl, 4-imidazoacetyl, or 4-imidazo-a, a dimethyl-acetyl; R² is selected from the group consisting of C₆-C₁₀ unbranched acyl, or is absent; R³ is selected from the group consisting of

-12-

Gly-OH or NH₂; and, Xaa is Lys or Arg, may be used in present invention.

More preferred compounds of SEQ ID NO:3 for use in the present invention are those in which Xaa is Arg and
5 R² is C₆-C₁₀ unbranched acyl.

Highly preferred compounds of SEQ ID NO:3 for use in the present invention are those in which Xaa is Arg, R² is C₆-C₁₀ unbranched acyl, and R³ is Gly-OH.

More highly preferred compounds of SEQ ID NO:3
10 for use in the present invention are those in which Xaa is Arg, R² is C₆-C₁₀ unbranched acyl, R³ is Gly-OH, and R¹ is 4-imidazopropionyl.

The most preferred compound of SEQ ID NO:3 for use in the present invention is that in which Xaa is Arg, R² is
15 C₈ unbranched acyl, R³ is Gly-OH, and R¹ is 4-imidazopropionyl.

The use of Val⁸-GLP-1(7-37)OH or a pharmaceutically-acceptable salt thereof, as claimed in US Patent Number 5,705,483, herein incorporated by reference,
20 in the present invention is highly preferred.

Methods for preparing the GLP-1, GLP-1 analogs, or GLP-1 derivatives useful in the present invention are well-known in the art and are easily within the grasp of ordinarily skilled protein chemists or biochemists. The
25 amino acid portion of the active compound used in the present invention, or a precursor thereto, can be made either by solid-phase synthetic chemistry, purification of GLP-1 molecules from natural sources, or recombinant DNA

-13-

technology. Routine synthetic organic techniques enable the alkylation and acylation of the GLP-1 derivatives.

The term "GLP-1 related compound" refers to any compound falling within the GLP-1, GLP-1 analog, or GLP-1
5 derivative definition.

The term "preservative" refers to a compound added to a pharmaceutical formulation to act as an anti-microbial agent. A parenteral formulation must meet guidelines for preservative effectiveness to be a commercially viable
10 multi-use product. Among preservatives known in the art as being effective and acceptable in parenteral formulations are benzalkonium chloride, benzethonium, chlorohexidine, phenol, m-cresol, benzyl alcohol, methylparaben, chlorobutanol, o-cresol, p-cresol, chlorocresol,
15 phenylmercuric nitrate, thimerosal, benzoic acid, and various mixtures thereof. See, e.g., Wallhauser, K., *Develop. Biol. Standard*, 24: 9-28 (Basel, S. Krager, 1974).

The term "buffer" or "pharmaceutically acceptable buffer" refers to a compound that is known to be safe for
20 use in protein formulations and that has the effect of controlling the pH of the formulation at the pH desired for the formulation. Pharmaceutically acceptable buffers for controlling pH at a moderately acid pH to a moderately basic pH include, for example, such compounds as phosphate,
25 acetate, citrate, TRIS, arginine, or histidine.

The term "isotonicity agent" refers to a compound that is tolerated physiologically and imparts a suitable tonicity to a formulation to prevent the net flow of water across the cell membrane. Compounds, such as glycerin, are
30 commonly used for such purposes at known concentrations.

-14-

group consisting of a peptide having the amino acid
sequence:

His-Ala-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-Ser-Tyr-
Leu-Glu-Gly-Gln-Ala-Ala-Lys-Glu-Phe-Ile-Ala-Trp-Leu-
5 Val-X (SEQ ID NO:2)

wherein X is selected from the group consisting of Lys and
Lys-Gly; and a derivative of said peptide, wherein said
peptide is selected from the group consisting of: a
pharmaceutically-acceptable acid addition salt of said
10 peptide; a pharmaceutically-acceptable carboxylate salt of
said peptide; a pharmaceutically-acceptable lower alkylester
of said peptide; and a pharmaceutically-acceptable amide of
said peptide selected from the group consisting of amide,
lower alkyl amide, and lower dialkyl amide.

15 Another preferred group of molecules for use in
the present invention consists of compounds disclosed in
U.S. Patent No. 5,512,549, herein incorporated by reference,
having the general formula:

20 R¹-Ala-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-Ser-Tyr-
Leu-Glu-Gly-Gln-Ala-Ala-Xaa-Glu-Phe-Ile-Ala-Trp-
Leu-Val-Lys-Gly-Arg-R³
R²

(SEQ ID NO:3)

25 and pharmaceutically-acceptable salts thereof, wherein R¹ is
selected from the group consisting of 4-imidazopropionyl, 4-

-15-

in the level of circulating insulin followed by a rapid fall in blood glucose levels. Different inhalation devices typically provide similar pharmacokinetics when similar particle sizes and similar levels of lung deposition are compared.

According to the invention, GLP-1 and GLP-1 analogs and derivatives can be delivered by any of a variety of inhalation devices known in the art for administration of a therapeutic agent by inhalation. These devices include metered dose inhalers, nebulizers, dry powder inhalers, sprayers, and the like. Preferably, GLP-1 and GLP-1 analogs and derivatives are delivered by a dry powder inhaler or a sprayer. There are a several desirable features of an inhalation device for administering GLP-1 and GLP-1 analogs and derivatives. For example, delivery by the inhalation device is advantageously reliable, reproducible, and accurate. The inhalation device should deliver small particles, e.g. less than about 10 μm mass median aerodynamic diameter (MMAD), preferably about 1-5 μm MMAD, for good respirability. Some specific examples of commercially available inhalation devices, or those in late stage development, suitable for the practice of this invention are TurbohalerTM (Astra), Rotahaler[®] (Glaxo), Diskus[®] (Glaxo), SpirosTM inhaler (Dura), devices being developed by Inhale Therapeutics, AERxTM (Aradigm), the Ultravent[®] nebulizer (Mallinckrodt), the Acorn II[®] nebulizer (Marquest Medical Products), the Ventolin[®] metered dose inhaler (Glaxo), the Spinhaler[®] powder inhaler (Fisons), or the like.

-16-

As those skilled in the art will recognize, the formulation of GLP-1 and GLP-1 analogs and derivatives, the quantity of the formulation delivered, and the duration of administration of a single dose depend on the type of inhalation device employed. For some aerosol delivery systems, such as nebulizers, the frequency of administration and length of time for which the system is activated will depend mainly on the concentration of the GLP-1 molecule in the aerosol. For example, shorter periods of administration can be used at higher concentrations of GLP-1 and GLP-1 analogs and derivatives in the nebulizer solution. Devices such as metered dose inhalers can produce higher aerosol concentrations, and can be operated for shorter periods to deliver the desired amount of GLP-1 and GLP-1 analogs and derivatives. Devices such as powder inhalers deliver active agent until a given charge of agent is expelled from the device. In this type of inhaler, the amount of GLP-1 and GLP-1 analogs and derivatives in a given quantity of the powder determines the dose delivered in a single administration.

The particle size of the GLP-1 and GLP-1 analogs and derivatives in the formulation delivered by the inhalation device is critical with respect to the ability of protein to deposit in the lungs, and preferably in the lower airways or alveoli. Preferably, the GLP-1 and GLP-1 analogs and derivatives is formulated so that at least about 10% of the peptide delivered is deposited in the lung, preferably about 10% to about 20%, or more. It is known that the maximum efficiency of pulmonary deposition for mouth breathing humans is obtained with particle sizes of about 2

-17-

μm to about 3 μm MMAD. When particle sizes are above about 5 μm MMAD, pulmonary deposition decreases substantially. Particle sizes below about 1 μm MMAD cause pulmonary deposition to decrease, and it becomes difficult to deliver particles with sufficient mass to be therapeutically effective. Thus, particles of GLP-1 and GLP-1 analogs and derivatives delivered by inhalation have a particle size preferably less than about 10 μm MMAD, more preferably in the range of about 1 μm to about 5 μm MMAD, and most preferably in the range of about 2 μm to about 3 μm MMAD. The formulation of GLP-1 and GLP-1 analogs and derivatives is selected to yield the desired particle size in the chosen inhalation device.

Advantageously for administration as a dry powder, GLP-1 and GLP-1 analogs and derivatives are prepared in a particulate form resulting in an emitted particle size less than about 10 μm MMAD, preferably about 1 to about 5 μm MMAD, and most preferably about 2 μm to about 3 μm MMAD. The preferred particle size is effective for delivery to the alveoli of the patient's lung. Preferably, the dry powder is largely composed of particles produced so that a majority of the particles have a size in the desired range. Advantageously, at least about 50% of the dry powder is made of particles having a diameter less than about 10 μm MMAD. Such formulations can be achieved by spray drying, milling, or critical point condensation of a solution containing the particular GLP-1 molecule and other desired ingredients. Other methods also suitable for generating particles useful in the current invention are known in the art.

-18-

The particles are usually separated from a dry powder formulation in a container and then transported into the lung of a patient via a carrier air stream. Typically, in current dry powder inhalers, the force for breaking up the solid is provided solely by the patient's inhalation. One suitable dry powder inhaler is the Turbohaler[™] manufactured by Astra (Södertälje, Sweden). In another type of inhaler, air flow generated by the patient's inhalation activates an impeller motor which deagglomerates the GLP-1 molecule particles. The Dura Spiros[™] inhaler is such a device.

Formulations of GLP-1 and GLP-1 analogs and derivatives for administration from a dry powder inhaler typically include a finely divided dry powder containing peptide, but the powder can also include a bulking agent, carrier, excipient, another additive, or the like. Additives can be included in a dry powder formulation of GLP-1 and GLP-1 analogs and derivatives, for example, to dilute the powder as required for delivery from the particular powder inhaler, to facilitate processing of the formulation, to provide advantageous powder properties to the formulation, to facilitate dispersion of the powder from the inhalation device, to stabilize the formulation (e.g., antioxidants or buffers), to provide taste to the formulation, or the like. Advantageously, the additive does not adversely affect the patient's airways. The GLP-1 and GLP-1 analogs and derivatives can be mixed with an additive at a molecular level or the solid formulation can include particles of the peptide mixed with or coated on particles of the additive. Typical additives include mono-, di-, and

-19-

polysaccharides; sugar alcohols and other polyols, such as, for example, lactose, glucose, raffinose, melezitose, lactitol, maltitol, trehalose, sucrose, mannitol, starch, or combinations thereof; surfactants, such as sorbitols, 5 diphosphatidyl choline, or lecithin; or the like. Typically an additive, such as a bulking agent, is present in an amount effective for a purpose described above, often at about 50% to about 90% by weight of the formulation. Additional agents known in the art for formulating proteins 10 can also be included in the formulation.

In another aspect of the invention a spray including GLP-1 and GLP-1 analogs and derivatives can be produced by forcing a suspension or solution of the peptide through a nozzle under pressure. The nozzle size and 15 configuration, the applied pressure, and the liquid feed rate can be chosen to achieve the desired output and particle size. An electro-spray can be produced, for example, by an electric field in connection with a capillary or nozzle feed. Advantageously, proplets of GLP-1 and GLP-1 20 analogs and derivatives delivered by a sprayer have an inhaled droplet size less than about 10 μm MMAD, preferably in the range of about 1 μm to about 5 μm MMAD, and most preferably about 2 μm to about 3 μm MMAD.

Formulations of GLP-1 and GLP-1 analogs and 25 derivatives suitable for use with a sprayer typically are about 1 mg to about 20 mg of the peptide per ml of solution. The formulation can include agents such as an excipient, a buffer, an isotonicity agent, a preservative, a surfactant, and metal cations. The formulation can also include an 30 excipient or agent to stabilize the peptide such as a

-20-

buffer, a reducing agent, a bulk protein, or a carbohydrate. Bulk proteins useful in formulating GLP-1 and GLP-1 analogs and derivatives include albumin, protamine, or the like. Typical carbohydrates useful in formulating GLP-1 and GLP-1
5 analogs and derivatives include sucrose, mannitol, lactose, trehalose, glucose, or the like. Formulations of GLP-1 and GLP-1 analogs and derivatives can also include a surfactant, which can reduce or prevent surface-induced aggregation of the peptide caused by atomization of the solution in forming
10 an aerosol. Various conventional surfactants can be employed, such as polyoxyethylene fatty acid esters and alcohols, and polyoxyethylene sorbitol fatty acid esters. Amounts will generally range between 0.001 and 4% by weight of the formulation. Other surfactants such as
15 diphosphatidyl choline or lecithin can also be used. Especially preferred surfactants for purposes of this invention are polyoxyethylene sorbitan monooleate, polysorbate 80, polysorbate 20, or the like. Additional agents known in the art for formulating proteins can also be
20 included in the formulation.

GLP-1 and GLP-1 analogs and derivatives can be administered by a nebulizer, such as jet nebulizer or an ultrasonic nebulizer. Typically, in a jet nebulizer, a compressed air source is used to create a high-velocity air
25 jet through an orifice. As the gas expands beyond the nozzle, a low-pressure region is created, which draws a solution of the peptide through a capillary tube connected to a liquid reservoir. The liquid stream from the capillary tube is sheared into unstable filaments and droplets as it
30 exits the tube, creating the aerosol. A range of

-21-

configurations, flow rates, and baffle types can be employed to achieve the desired performance characteristics from a given jet nebulizer. In an ultrasonic nebulizer, high-frequency electrical energy is used to create vibrational, mechanical energy, typically employing a piezoelectric transducer. This energy is transmitted to the peptide formulation either directly or through a coupling fluid, creating an aerosol. Advantageously, droplets of GLP-1 and GLP-1 analogs and derivatives delivered by a nebulizer have a particle size less than about 10 μm MMAD, preferably in the range of about 1 μm to about 5 μm MMAD, and most preferably about 2 μm to about 3 μm MMAD.

Formulations of GLP-1 and GLP-1 analogs and derivatives suitable for use with a nebulizer, either jet or ultrasonic, typically include an aqueous solution of the peptide at a concentration of about 1 mg to about 20 mg per ml of solution. The formulation can include agents such as an excipient, a buffer, an isotonicity agent, a preservative, a surfactant, and a divalent metal cation. The formulation can also include an excipient or agent to stabilize the peptide, such as a buffer, a reducing agent, a bulk protein, or a carbohydrate. Bulk proteins useful in formulating GLP-1 and GLP-1 analogs and derivatives include albumin, protamine, or the like. Typical carbohydrates useful in formulating GLP-1 related proteins include sucrose, mannitol, lactose, trehalose, glucose, or the like. Formulations of the GLP-1 and GLP-1 analogs and derivatives can also include a surfactant, which can reduce or prevent surface-induced aggregation of the peptide caused by atomization of the solution in forming an aerosol. Various

-22-

conventional surfactants can be employed, such as polyoxyethylene fatty acid esters and alcohols, and polyoxyethylene sorbital fatty acid esters. Amounts will generally range between 0.001 and 4% by weight of the formulation. Other surfactants such as phosphatidyl choline or lethicin can also be used. Especially preferred surfactants for purposes of this invention are polyoxyethylene sorbitan monooleate, polysorbate 80, polysorbate 20, or the like. Additional agents known in the art for formulation of a protein such as GLP-1 related molecules can also be included in the formulation.

Another aspect of the invention involves a metered dose inhaler (MDI). In this embodiment, a propellant, GLP-1 and GLP-1 analogs and derivatives, and any excipients or other additives are contained in a canister as a mixture including a liquefied compressed gas. Actuation of the metering valve releases the mixture as an aerosol, preferably containing inhaled particles in the size range of less than about 10 μm MMAD, preferably about 1 μm to about 5 μm MMAD, and most preferably about 2 μm to about 3 μm MMAD. The desired aerosol particle size can be obtained by employing a formulation of GLP-1 and GLP-1 analogs and derivatives produced by various methods known to those of skill in the art, including jet-milling, spray drying, critical point condensation, or the like. Preferred metered dose inhalers include those manufactured by 3M or Glaxo and employing a hydrofluorocarbon propellant.

Formulations of GLP-1 and GLP-1 analogs and derivatives for use with a metered-dose inhaler device will generally include a finely divided powder containing peptide

-23-

as a suspension in a non aqueous medium, for example, suspended in a propellant with the aid of a surfactant. The propellant may be any conventional material employed for this purpose, such as chlorofluorocarbon, a

5 hydrochlorofluorocarbon, a hydrofluorocarbon, or a hydrocarbon, including trichlorofluoromethane, dichlorodifluoromethane, dichlorotetrafluoroethanol and 1,1,1,2-tetrafluoroethane, HFA-134a (hydrofluroalkane-134a), HFA-227 (hydrofluroalkane-227), or the like. Preferably the

10 propellant is a hydrofluorocarbon. The surfactant can be chosen to stabilize the GLP-1 molecule as a suspension in the propellant, to protect the active agent against chemical degradation, and the like. Suitable surfactants include sorbitan trioleate, soya lecithin, oleic acid, or the like.

15 In some cases solution aerosols are preferred using solvents such as ethanol. Other surfactants such as diphosphatidyl choline or lethicin can also be used. Additional agents known in the art for formulating proteins can also be included in the formulation.

20 The present invention also relates to a pharmaceutical composition or formulation including GLP-1 and GLP-1 analogs and derivatives and suitable for administration by inhalation. According to the invention, GLP-1 and GLP-1 analogs and derivatives can be used for

25 manufacturing a formulation or medicament suitable for administration by inhalation. The invention also relates to methods for manufacturing formulations including GLP-1 related molecules in a form that is suitable for administration by inhalation. For example, a dry powder

30 formulation can be manufactured in several ways, using

-24-

conventional techniques. Particles in the size range appropriate for maximal deposition in the lower respiratory tract can be made by micronizing, milling, spray drying, or the like. And a liquid formulation can be manufactured by
5 dissolving the peptide in a suitable solvent, such as water, at an appropriate pH, including buffers or other excipients.

The present invention may be better understood with reference to the following examples. These examples are intended to be representative of specific embodiments of
10 the invention, and are not intended as limiting the scope of the invention.

EXAMPLES

Serum Pharmacokinetics of Val⁸-GLP-1 in 15 Beagle Dogs Following Pulmonary Administration

The GLP-1 analog, Val⁸-GLP-1(7-37)OH (SEQ ID NO:4) was prepared in *E coli* using conventional recombinant DNA techniques and purified to homogeneity.

NH₂-His-Val-Glu-Gly-Thr-Phe-Thr-Ser-Asp-Val-Ser-Ser-Tyr-Leu-
20 Glu-Gly-Gln-Ala-Ala-Lys-Glu-Phe-Ile-Ala-Trp-Leu-Val-Lys-Gly-Arg-Gly-OH

(SEQ ID NO: 4)

One group of 6 female beagle dogs was exposed to inhaled Val⁸-GLP-1 for 15 minutes at an average aerosol
25 concentration of 77.2 µg/L generated from a solution of Val⁸-GLP-1 in sterile water. These animals were dosed with 100 µg/kg via subcutaneous administration approximately 1-week after inhalation exposures. Tidal volume, breathing rate, and minute volume were monitored prior to and
30 throughout the exposure period. Blood was collected for

-25-

analysis of plasma levels of Val⁸-GLP-1 at various time points following inhalation and subcutaneous administration. Bronchoalveolar lavage (BAL) fluid was collected approximately 4 hours postexposure and analyzed for LDH, total protein, cell counts, and white cell differentials.

The delivery of Val⁸-GLP-1 was well tolerated with an inhaled dose of 1198 µg/kg and an estimated deposited lung dose of 240 µg/kg. Subcutaneous administration of 100 µg/kg was also well tolerated by all animals. Inhalation and subcutaneous administration of Val⁸-GLP-1 was delivered using formulated and formulated material.

There were no treatment-related clinical observations; body weights were not adversely affected by Val⁸-GLP-1. Only minor lung effects were observed. Increases in both tidal volume and minute volume were observed during the 15 minute inhalation exposures but the data were highly variable. No significant changes were observed for LDH, red blood cell counts, white blood cell counts, neutrophils, lymphocytes, eosinophils, epithelial cells, macrophages, basophils, or monocytes. There was a mild increase in total protein following aerosol delivery of Val⁸-GLP-1.

Results from this study demonstrated that there was good bioavailability of Val⁸-GLP-1 (40%, based on AUC) delivered to the lungs of beagle dogs by inhalation relative to subcutaneous administration. Val⁸-GLP-1 was well tolerated for up to 15 minutes with minimal effects on the lungs at an average inhaled dose of 1198 µg/kg, which was a no-observed-adverse-effects level (NOAEL) in this study.

-26-

Preparation of Dose Solutions

Solutions of Val⁸-GLP-1 were prepared on the days of dosing at concentrations of 0.5 mg/ml or 8 mg/ml in sterile water for subcutaneous administration and pulmonary administration, respectively. An additional solution of Val⁸-GLP-1 (8 mg/ml) was prepared at the end of live phase in order to determine the particle size distribution of aerosolized Val⁸-GLP-1. The solutions were filtered through a low protein binding 0.22 micron canister filter. The pH of the solution was adjusted with sodium hydroxide solution to 7.47.

Test Animals

Six females Beagle dogs (Marshall Farms, North Rose, NY) were used in this study. Each animal was uniquely identified by a five-digit animal number and a seven-digit tattoo (located on inner ear) number recorded on their cage card. All of the animals were acclimated to the restraint slings prior to beginning the study. The weight range of the animals at the start of the study was 8.4 to 11.1 kg. The age of the animals at the start of the study was 33 to 37 weeks.

Test Animal Housing and Care

Animals were pair housed in stainless steel cages except on the days of exposure. Each animal was individually housed on the day of exposure in order to monitor their feeding regimen. Rooms were thermostatically set to maintain a temperature of 70°F and maintain the actual temperature within $\pm 8^\circ\text{F}$ from that set point. The environmental control system is designed to maintain a

-27-

relative humidity of 20% and a maximum of 80%. Light was on a 12-hour cycle, with lights on between 0600 and 1800 hours. Subsequently, lights were off between 1800 and 0600 hours except when blood samples were collected. Animals were fed
5 once daily with Hill's Science Diet. Animals were fasted for approximately 12-hours prior to exposure. Tap water was provided *ad libitum* except during exposures.

Treatment Groups and Study Duration

All 6 dogs were exposed for 15 minutes to
10 aerosolized Val⁸-GLP-1. The targeted deposited lung dose for Val⁸-GLP-1 exposures was 200 µg/kg of body weight. Approximately 7 days after pulmonary administration of Val⁸-GLP-1, all dogs were dosed with 100 µg/kg Val⁸-GLP-1 of body weight via subcutaneous administration.

15

Exposure System

The dogs were tested while standing in restraint slings. Two layers of 0.03 inch latex sheets were placed around the animals' necks to form a nonrestrictive airtight
20 seal. A custom built, 11-L head-dome, similar to that described by Allen et al. (J Appl Toxicol 1995; 15:13-17) was placed over the dogs' heads and secured to the sling. Airflow was exhaust driven via a transvector located on the exhaust side of the dome. Because the helmet was airtight
25 and the neck was sealed, this constituted a head-only exposure system. The total flow rate through the dome was approximately 7.5 L/min.

Aerosol Generation

The aerosols were generated using a Respigard II
30 nebulizer with an input of approximately 6.5 L/min. The

-28-

output from the generator flowed directly into the head-dome.

Atmospheric Concentration Sampling

All sampling for total gravimetric concentrations was performed with in-line filter holders containing type A/E glass fiber filters (Gelman Instruments Co., Ann Arbor, MI). Filters sampling from the chamber was carried out at a nominal sampling rate of 1 L/min, calibrated with a portable mass airflow calibrator (Model 830, Sierra Instruments, Carmel Valley, CA). Sampling duration was 15 minutes. Particle size analysis was performed with a Sierra Model 218K Ambient Cascade Impactor (Anderson Samplers, Inc, Atlanta, GA). Cascade sampling from the chamber was carried out at a nominal sampling rate of 3 L/min, calibrated with a portable mass airflow calibrator (Model 830, Sierra Instruments, Carmel Valley, CA). Sampling duration was 31 minutes. Filters were allowed to dry for approximately 30-minutes before being re-weighed.

Dose Determination

The dose of Val⁸-GLP-1 inhaled during a 15-minute exposure was estimated as follows: the mean minute volume (mL) during the 15 minute exposure was multiplied by the exposure duration to yield the total air breathed (L) during the inhalation exposure. This value was multiplied by the aerosol concentration ($\mu\text{g/L}$) to determine total dose (mg). The inhaled dose ($\mu\text{g/kg}$) was calculated by dividing the total dose (μg) by the animal's body weight (kg).

The dose of Val⁸-GLP-1 deposited in the lungs was estimated as follows: inhaled dose ($\mu\text{g/kg}$) was multiplied

-29-

by 20 percent to yield estimated deposited lung dose
($\mu\text{g}/\text{kg}$). Aerosols with a MMEAD ranging from 1-2 μm MMAD
have been shown to deposit in the lung with approximately a
20 percent efficiency (Schlesinger RB. 1985. Comparative
5 deposition of inhaled aerosols in experimental animals and
humans: a review. J Toxicol Environ Health 15:197-214).

Pulmonary Function

All animals were weighed on Days -5, 0, and 7.
Breathing patterns (tidal volume, breathing frequency, and
10 minute volume) were monitored using a size '0'
pneumotachograph connected to a port on the head-dome. The
signals were collected on a personal computer using the
Buxco XA Data Acquisition System (Buxco Electronics, Inc,
Sharon, CT). At least 15 minutes of preexposure data were
15 collected before the exposures began, followed by data
collection throughout the 15-minute exposure periods. All
data were analyzed as 5-minute averages.

Bronchoalveolar Lavage

Bronchoalveolar lavage (BAL) was performed
20 approximately 4 hours after each dosing regimen. The
animals were anesthetized with an intravenous injection of
2% Brevital prior to the BAL procedure. Bronchoalveolar
lavage was performed using a pediatric fiberoptic
bronchoscope (Olympus, model BF, Type 3C10, Lake Success,
25 NY). The bronchoscope tip was wedged in a 5th to 7th
generation airway of a lower lobe. The BALs were alternated
between the right and left lung lobes. Two 10-mL aliquots
were instilled and gently suctioned out. Aliquots of the
recovered lavage fluid were used to determine total white

-30-

cell counts, total red cell counts, white cell differentials, total protein, and lactate dehydrogenase.

Cell Counts and Differentials

A complete blood count was done on the
5 unconcentrated BAL fluid using a Technicon H1 System (Technicon Instruments Corporation). Cell differentials for the BAL were done by microscopic evaluation of 200 Wright stained cells.

Blood Collection

10 For analysis of plasma pharmacokinetics of Val⁸-GLP-1, approximately 2 to 3 mL of blood were collected in EDTA vacutainer tubes from the cephalic or jugular vein prior to exposure and at 0.25, 0.5, 1, 2, 3, 4, 6, 8, and 12 hours postexposure. To obtain the plasma, each tube was
15 centrifuged at approximately 3000 rpm for 10 minutes at 10°C. The plasma samples were stored at -70 °C until sent for assay.

Exposure Concentration/Particle Size and Lung Dose

The average exposure concentration for each dog
20 exposed to Val⁸-GLP-1 ranged from 64.0 to 101.3 µg/L. The mean (\pm SD) concentration for all animals was 77.2 ± 16.9 µg/L. Particle size was measured as a mass median equivalent aerodynamic diameter (MMEAD) of 0.91 µm with a geometric standard deviation (GSD) of 2.37.

25 The average dose of Val⁸-GLP-1 deposited in lungs of 6 dogs treated with Val⁸-GLP-1 was calculated as 240 ± 42 µg/kg (mean \pm SD). The mean inhaled dose (that which entered the respiratory tract neglecting deposition) was 1198 ± 208 µg/kg (mean \pm SD). Individual animal data are
30 shown in the table below.

-31-

Estimated Lung Dose for Animals Exposed
to Aerosols of Val⁸-GLP-1

Animal Number	Aerosol Conc. (µg/L)	Average Minute Volume (ml)	Inhaled Lung Dose (µg/kg)	Deposited Lung Dose (µg/kg)
27682	0.07717*	8406	1060	212
27684	0.07717*	9563	1030	206
27685	0.10133	7260	1350	270
27686	0.06400	12733	1360	272
27687	0.06733	9214	950	190
27689	0.07600	11890	1400	288

- 5 **Aerosol concentration was not determined for animals 27682 and 27684, therefore the mean aerosol concentration was used to calculate an estimated inhaled lung dose (µg/kg).*

No significant changes in body weight were
 10 observed during the treatment phase. The initial (Day -5)
 and final (Day 7) body weights were 9.5 ± 0.9 (mean \pm SD;
 n=6) and 9.7 ± 0.9 kg, respectively. No marked changes in
 breathing frequency was observed. Slight increases in tidal
 volume and minute volume were measured but the data was
 15 highly variable potentially due to the short acclimation
 period prior to study initiation. Individual animal data
 are shown in the table below.

- 32 -

Changes in Pulmonary Function During
Exposure to Val⁸-GLP-1

Animal	Tidal Volume (mL)			Breathing Frequency (bpm)			Minute Volume (mL/min)		
Number	5*	10*	15*	5*	10*	15*	5*	10*	15*
27682	119.0	152.1	105.1	74	86.0	95.7	7265	9360	8592
27684	104.6	99.4	100.8	119.2	118.2	108.7	10300	9117	9272
27685	165.9	178.5	209.6	58.1	49.4	42.5	7938	7465	6377
27686	325.1	586.2	374.4	56.3	33.8	59.4	12200	13300	12700
27687	245.9	221.6	183.1	39.5	45.5	50.5	9288	9573	8781
27689	184.5	367.6	454.0	39.8	46.8	33.3	7271	13400	15000
Mean	190.8	267.6	237.8	64.6	63.3	65.0	9044	10369	10120
Stdev	82.8	180.7	145.3	29.8	32.2	30.3	1956	2426.1	3141
Mean	239.0	216.8	207.6	45.9	51.4	64.4**	8448	8744.8	8223**
(baseline)			**						
Stdev	161.4	137.2	209.5	12	10.2	32.3	2614	2614	1566
(baseline)									

*minutes during exposure (values represent average over 5 minutes)

**mean calculated from a n=5 (instead of n=6) because no value was recorded for animal

#27689 at 15 minutes preexposure

Comparison of Pharmacokinetics Following Subcutaneous and
Pulmonary Administration of Val⁸-GLP-1

5

Delivery	C _{max}	T _{max}	AUC _{0-t(1)}	T1/2(α)
Route	(ng/mL)	(h)	(ng*h/mL)	(hours)
Subcutaneous	10.53 ± 1.06	0.71 ± 0.14	36.37 ± 2.18	1.26 ± 0.11
Inhalation	8.66 ± 0.90	1.54 ± 0.59	35.21 ± 5.91	1.19 ± 0.11

*All values are reported as mean ± SEM (standard error of the mean).

(1) AUC_{0-t}=area under the plasma concentration curve from time 0 to t, where t=12 hours postdose

10 Plasma concentrations of immunoreactive Val⁸-GLP-1 (Table 3) were measured by a competitive radioimmunoassay

-33-

(RIA). The absorption of Val⁸-GLP-1 via both delivery routes appeared to be rapid, reaching substantial plasma concentrations at 15 minutes postdose. The plasma time profiles were similar for the subcutaneous injections and inhalation. The average Tmax value for inhalation was greater than that for subcutaneous injection. Also, the elevated plasma Val⁸-GLP-1 concentration (close to Cmax) achieved by inhalation appeared to remain near that level for a longer period of time than following subcutaneous injection.

Based on the average AUC values, the bioavailability of inhaled Val⁸-GLP-1 (averaged inhaled dose of 1198 µg/kg) relative to subcutaneous injection (100 µg/kg) was approximately 7.7%. On the basis of deposited lung dose, estimated as 240 µg/kg, the bioavailability relative to subcutaneous injection was 40%.

-34-

WE CLAIM:

1. A method of administering a glucagon-like peptide-1 (GLP-1) molecule comprising, administering an effective amount of a GLP-1 molecule selected from the group
5 consisting of GLP-1, GLP-1 analogs, or GLP-1 derivatives to a patient in need thereof by pulmonary means.

2. The method of Claim 1, wherein the GLP-1 molecule is delivered to lower airway of the patient.

10

3. The method of Claim 2, wherein the GLP-1 molecule is deposited in the alveoli.

4. The method of Claim 1, wherein the GLP-1 molecule
15 is inhaled through the mouth of the patient.

5. The method of Claim 1, wherein the GLP-1 molecule is administered as a pharmaceutical formulation comprising the GLP-1 molecule in a pharmaceutically acceptable carrier.

20

6. The method of Claim 5, wherein the formulation is selected from the group consisting of a solution in an aqueous medium and a suspension in a non-aqueous medium.

25 7. The method of Claim 6, wherein the formulation is administered as an aerosol.

8. The method of Claim 5, wherein the formulation is in the form of a dry powder.

30

-35-

9. The method of Claim 5, wherein the GLP-1 molecule has a particle size of less than about 10 microns MMAD.

10. The method of Claim 9, wherein the GLP-1 molecule has a particle size of about 1 to about 5 microns MMAD.

11. The method of Claim 10, wherein the GLP-1 molecule has a particle size of about 2 to about 3 microns MMAD.

12. The method of Claim 1, wherein at least about 10% of the GLP-1 molecule delivered is deposited in the lung.

13. The method of Claim 1, wherein the GLP-1 molecule is delivered from an inhalation device suitable for pulmonary administration and capable of depositing the GLP-1 molecule in the lungs of the patient.

14. The method of Claim 13, wherein the device is selected from the group consisting of a nebulizer, a metered-dose inhaler, a dry powder inhaler, and a sprayer.

15. The method of Claim 14, wherein the device is a dry powder inhaler.

16. The method of Claim 1 wherein the GLP-1 molecule is selected from the group consisting of GLP-1 analogs and GLP-1 derivatives.

17. The method of Claim 16 wherein the GLP-1 molecule is a GLP-1 analog.

-36-

18. The method of Claim 17 wherein the GLP-1 analog is selected from the group consisting of Val⁸-GLP-1(7-37)OH, Gly⁸-GLP-1(7-37)OH, and Asp⁸-GLP-1(7-37)OH.

5

19. The method of Claim 18, wherein the GLP-1 analog is Val⁸-GLP-1(7-37)OH.

20. The method of Claim 18, wherein the GLP-1 analog is Gly⁸-GLP-1(7-37)OH.

10

21. A method for treating diabetes comprising, administering an effective dose of a GLP-1 molecule to a patient in need thereof by pulmonary delivery.

15

22. The method of Claim 21, wherein the GLP-1 molecule is administered as a pharmaceutical formulation comprising the GLP-1 molecule in a pharmaceutically acceptable carrier.

20

23. The method of Claim 21, wherein the GLP-1 molecule is Val⁸-GLP-1(7-37)OH.

24. The method of Claim 21, wherein the GLP-1 molecule is Gly⁸-GLP-1(7-37)OH.

25

25. The method of Claim 21, wherein the GLP-1 molecule is delivered from an inhalation device suitable for pulmonary administration and capable of depositing the GLP-1 molecule in the lungs of the patient.

30

-37-

26. The method of Claim 25, wherein the device is a sprayer or a dry powder inhaler.

27. The method of Claim 25, wherein an actuation of
5 the device administers about 40 µg to about 4,000 µg of a GLP-1 molecule.

28. The method of Claim 25, wherein an actuation of
the device administers about 80 µg to about 2,000 µg of a
10 GLP-1 molecule.

29. The method of Claim 25, wherein an actuation of
the device administers about 160 µg to about 1,000 µg of a
GLP-1 molecule.

15

30. The method of Claim 25, wherein an actuation of
the device administers about 320 µg to about 500 µg of a
GLP-1 molecule.

20 31. A method for treating hyperglycemia comprising,
administering an effective dose of a GLP-1 molecule to a
patient in need thereof by pulmonary means.

32. The method of Claim 31, wherein the GLP-1 molecule
25 is administered as a pharmaceutical formulation comprising
the GLP-1 molecule in a pharmaceutically acceptable carrier.

33. The method of Claim 31, wherein the GLP-1 molecule
is Val⁸-GLP-1(7-37)OH.

30

-38-

34. The method of Claim 31, wherein the GLP-1 molecule is Gly⁸-GLP-1(7-37)OH.

35. The method of Claim 31, wherein the GLP-1 molecule
5 is delivered from an inhalation device suitable for pulmonary administration and capable of depositing the GLP-1 molecule in the lungs of the patient.

36. The method of Claim 35, wherein the device is
10 selected from the group consisting of a sprayer and a dry powder inhaler.

37. The method of Claim 35, wherein an actuation of the device administers about 40 µg to about 4,000 µg of GLP-
15 1 molecule.

38. The method of Claim 35, wherein an actuation of the device administers about 80 µg to about 2,000 µg of the GLP-1 molecule.

20

39. The method of Claim 35, wherein an actuation of the device administers about 160 µg to about 1,000 µg of GLP-1 molecule.

25 40. The method of Claim 35, wherein an actuation of the device administers about 320 µg to about 500 µg of the GLP-1 molecule.

SEQUENCE LISTING

<110> Eli Lilly and Company

<120> METHOD FOR ADMINISTERING INSULINOTROPIC PEPTIDES

<130> X-12013

<140>

<141>

<150> US 60/098273

<151> 1998-08-28

<150> US 60/100012

<151> 1998-09-11

<160> 4

<170> PatentIn Ver. 2.0

<210> 1

<211> 29

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: synthetic
sequence

<220>

<223> Xaa at position 1 is selected from the group
consisting of Ala, Gly, Val, Thr, Ile and
alpha-methyl-Ala.

<220>

<223> Xaa at position 14 is selected from the group
consisting of Glu, Gln, Ala, Thr, Ser and Gly.

<220>

<223> Xaa at position 20 is selected from the group
consisting of Glu, Gln, Ala, Thr, Ser and Gly.

<400> 1

Xaa	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Val	Ser	Ser	Tyr	Leu	Xaa	Gly	Gln
1				5				10					15		

Ala Ala Lys Xaa Phe Ile Ala Trp Leu Val Lys Gly Arg

20

25

<210> 2

<211> 28

<212> PRT

<213> Homo sapiens

<220>

<223> Description of Artificial Sequence: synthetic
sequence

<220>

<223> Xaa at position 28 is selected from the group
consisting of Lys and Lys-Gly.

<400> 2

His	Ala	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Val	Ser	Ser	Tyr	Leu	Glu	Gly
1				5					10					15	

Gln	Ala	Ala	Lys	Glu	Phe	Ile	Ala	Trp	Leu	Val	Xaa
			20					25			

<210> 3

<211> 29

<212> PRT

<213> Artificial Sequence

<220>

<223> Xaa at position 19 is selected from the group
consisting of Lys and Arg.

<220>

<223> Description of Artificial Sequence: synthetic
sequence

<400> 3

Ala	Glu	Gly	Thr	Phe	Thr	Ser	Asp	Val	Ser	Ser	Tyr	Leu	Glu	Gly	Gln
1				5					10					15	

Ala	Ala	Xaa	Glu	Phe	Ile	Ala	Trp	Leu	Val	Lys	Gly	Arg
			20					25				

<210> 4

<211> 31

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: synthetic or
semi-synthetic sequence

<400> 4

His Val Glu Gly Thr Phe Thr Ser Asp Val Ser Ser Tyr Leu Glu Gly
1 5 10 15

Gln Ala Ala Lys Glu Phe Ile Ala Trp Leu Val Lys Gly Arg Gly
20 25 30

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/19348

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 38/16

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/12, 530/324

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WEST, CAS Online

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,545,618 A (BUCKLEY et al.) 13 August 1996, see entire document.	1-40
A	US 5,512,549 A (CHEN et al.) 30 April 1996, see entire document.	1-40

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

29 OCTOBER 1999

Date of mailing of the international search report

10 NOV 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DAVID LUKTON

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/19348

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

514/12